



Proceedings of the Ninth Prague Topological Symposium
Contributed papers from the symposium held in
Prague, Czech Republic, August 19–25, 2001

TRANSFINITE SEQUENCES OF CONTINUOUS AND BAIRE 1 FUNCTIONS ON SEPARABLE METRIC SPACES

MÁRTON ELEKES

ABSTRACT. We investigate the existence of well-ordered sequences of Baire 1 functions on separable metric spaces.

Any set $\mathcal{F}$ of real valued functions defined on an arbitrary set X is partially ordered by the pointwise order, that is $f \leq g$ iff $f(x) \leq g(x)$ for all $x \in X$. In other words put $f < g$ iff $f(x) \leq g(x)$ for all $x \in X$ and $f(x) \neq g(x)$ for at least one $x \in X$. Our aim will be to investigate the possible length of the increasing or decreasing well-ordered sequences of functions in $\mathcal{F}$ with respect to this order.

A classical theorem of Kuratowski asserts, that if $\mathcal{F}$ is the set of continuous or Baire 1 functions defined on a Polish space X , then there exists a monotone sequence of length ξ in $\mathcal{F}$ iff $\xi < \omega_1$ (see [2, §24. III.2']). Moreover, P. Komjáth proved in [1] that the corresponding question concerning Baire α functions for $2 \leq \alpha < \omega_1$ is independent of *ZFC*.

In the present paper we investigate what happens if we drop the condition of completeness and replace the Polish space X by a separable metric space.

Our main results are the following. Let $d(X)$ denote the density of a space X .

Theorem. *Let (X, ρ) be a metric space. Then there exists a well-ordered sequence of length ξ of continuous real-valued functions defined on X iff $\xi < d(X)^+$.*

Corollary. *A metric space is separable iff every well-ordered sequence of continuous functions defined on it is countable.*

Theorem. *There exists a separable metric space on which there exists a well-ordered sequence of length ω_1 of Baire 1 functions.*

2000 *Mathematics Subject Classification.* 26A21.

Key words and phrases. Baire 1 function, well-ordered sequence, metric spaces.

This is a research announcement. A complete article, written jointly with Kenneth Kunen, will be published elsewhere.

Márton Elekes, *Transfinite sequences of continuous and Baire 1 functions on separable metric spaces*, Proceedings of the Ninth Prague Topological Symposium, (Prague, 2001), pp. 91–92, Topology Atlas, Toronto, 2002.

Theorem. *The following statement: ‘There exists a separable metric space on which there exists a well-ordered sequence of length ω_2 of Baire 1 functions’ is independent of $ZFC + \neg CH$.*

Remark. During and after the conference Kenneth Kunen answered one of my questions, and also improved some of the results and proofs. These results will appear in a forthcoming joint paper.

REFERENCES

- [1] Péter Komjáth, *Ordered families of Baire-2-functions*, Real Anal. Exchange **15** (1989/90), no. 2, 442–444. MR **91d**:26007
- [2] K. Kuratowski, *Topology. Vol. I*, Academic Press, New York, 1966. MR 36 #840

DEPARTMENT OF ANALYSIS, EÖTVÖS LORÁND UNIVERSITY, BUDAPEST, PÁZMÁNY
PÉTER SÉTÁNY 1/C, 1117, HUNGARY
E-mail address: `emarci@cs.elte.hu`